

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM
maintained by the
CALIFORNIA DEPARTMENT OF FISH AND GAME
and supported by the
CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP
Database Version 8.1 (2005)

B469 Summer Tanager *Piranga rubra*
Family: Thraupidae Order: Passeriformes Class: Aves

Written by: D. Dobkin
Reviewed by: L. Mewaldt
Edited by: R. Duke, S. Granholm

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

An uncommon (formerly common) summer resident and breeder in desert riparian habitat along lower Colorado River; also occurs very locally elsewhere in southern California deserts. Found in additional desert and other localities in migration. It is a rare but regular migrant and winter visitor along the coast, mostly from Los Angeles Co. southward, scattered records occur in northern California. Breeds in mature, desert riparian habitat dominated by cottonwoods and willows (Grinnell and Miller 1944, McCaskie et al. 1979, 1988, Garrett and Dunn 1981).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Eats insects, spiders, and small fruits. Gleans from foliage and bark, and hawks flying insects. Eats many bees and wasps; often takes larvae from hives and nests (Bent 1958).

Cover: Cottonwoods and willows, especially older, dense stands along rivers and streams, provide nesting, feeding, and other cover.

Reproduction: Builds nest 3-6 m (10-20 ft) above ground on a large, horizontal limb of a cottonwood or willow or other riparian deciduous trees.

Water: No additional information found.

Pattern: Frequents cottonwood-willow associations of riparian habitats for breeding, feeding, cover, and other activities.

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity.

Seasonal Movements/Migration: Arrives on summer breeding grounds in April and usually departs by September. Transients occur elsewhere in interior mostly in May and June and September into November. Occurs along coast rarely but regularly from September to March and May to June.

Home Range: No data found.

Territory: In deciduous forest and edge habitat in Kansas, Fitch and Fitch (1955) estimated territory at 1.7 ha (4.2 ac). May hold a feeding territory in winter (Ehrlich et al. 1988).

Reproduction: Breeding season lasts from May into early August with a peak in June. Clutch size 3-5, usually 4 (Bent 1958). Incubation apparently about 12 days (Terres 1980). Altricial young tended by both parents.

Niche: A California Species of Special Concern. Much less common now than in the 1940s, primarily because of loss and fragmentation of mature cottonwood and willow stands, especially along the Colorado River (Grinnell and Miller 1944, Remsen 1978).

REFERENCES

- Bent, A. C. 1958. Life histories of North American Blackbirds, orioles, tanagers, and allies. U.S. Natl. Mus. Bull. 211. 549pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. The birder's handbook. Simon and Schuster, New York. 785pp.
- Fitch, H. S., and V. R. Fitch. 1955. Observations on the summer tanager in northeastern Kansas. Wilson Bull. 67:45-54.
- Garrett, K., and J. Dunn. 1981. Birds of southern California. Los Angeles Audubon Soc. 408pp.
- Grinnell, J., and A. H. Miller. 1944. The distribution of the birds of California. Pac. Coast Avifauna No. 27. 608pp.
- Harrison, C. 1978. A field guide to the nests, eggs and nestlings of North American birds. W. Collins Sons and Co., Cleveland, OH. 416pp.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1979. Birds of northern California, an annotated field list. 2nd ed. Golden Gate Audubon Soc., Berkeley. 84pp.
- McCaskie, G., P. De Benedictis, R. Erickson, and J. Morlan. 1988. Birds of northern California, an annotated field list. 2nd ed. Golden Gate Audubon Soc., Berkeley. Reprinted with suppl. 108pp.
- Remsen, J. V., Jr. 1978. Bird species of special concern in California. Calif. Dept. of Fish and Game, Sacramento. Wildl. Manage. Admin. Rep. No. 78-1. 54pp.
- Terres, J. K. 1980. The Audubon Society encyclopedia of North American birds. A. Knopf, New York. 1100pp.